

www.danielmyermd.com

Dr. Daniel Myer's Post-operative Instructions for Elbow / Wrist / Hand

Please call our office with questions: North Coast / St. Thomas Hospital (330) 535-3396

Post-operative follow-up:

Office:

Date / Time:

Medications / Diet

- 1. Eat only light, non-greasy foods today.
- 2. Take pain medications with food.
- 3. While taking pain medications, you may not operate a vehicle, heavy machinery or appliances.
- 4. While taking pain medications, you may not drink alcoholic beverages.
- 5. While taking pain medications, you may not make critical decisions or sign legal papers.
- 6. If you have any reactions to your medicines, stop taking them and call my office immediately.
- 7. Please keep in mind that itching is a very common side effect of narcotic pain medicines, and if not allergic, over the counter Benadryl (diphenhydramine) may be used as directed.
- 8. Please keep in mind that constipation is a very common side effect of narcotic pain medicines. We recommend that patients take precautions to prevent constipation:
 - a. Drink plenty of water (6-8 glasses of 8 oz. a day).
 - b. Avoid alcohol and excessive caffeine.
 - c. Eat plenty of fiber (fruits, vegetables and whole grains).
 - d. Take an over the counter stool softener: Colace (docusate) or Dulcolax (bisacodyl).

The following medicines: ______ were sent electronically to ______

The following medicines were printed:

Yes / No Please take a daily 81mg Aspirin - to help prevent blood clots

Activity / Exercise

1. Range of Motion:

You may move the elbow / wrist / hand as much as the dressing will allow (OK to remove sling when nerve block wears off)

You are in a splint – range of motion is limited. Please keep splint on at all times until follow-up. As long as the splint does not limit your motion, please move your fingers and wrist as much as possible to minimize swelling.

Yes / No Physical Therapy script printed today. Please call to get appointment setup as discussed.

- 2. It is important to keep ice applied as long as pain or swelling persists. Do not apply ice directly to skin or allow water to leak on your dressing.
- 3. It is important to keep your extremity elevated as much as possible (on pillows) to help with swelling.

Dressing / Wound Care

- 1. Please keep dressing dry.
 - You can remove your dressing 48 hours after surgery. At that time, you can begin showering and get your incision/s wet. Simply dry the incision after a shower and cover with a dry dressing or Band-Aid.
 - Your splint must stay on until next follow-up. Please keep covered at all times with bathing.
- 2. Even after changing the dressing, it is normal to have some clear or bloody drainage from your incision/s.
- 3. Please absolutely NO BATHS, HOT TUBS or POOLS.

Emergency / Follow-up

- 1. Please notify my office if you develop any fever (101° or above), unexpected warmth, redness or swelling, or severe increased pain in your shoulder.
- 2. Please call if your fingers become suddenly cold, purple, prolonged numbress or there is excessive bleeding.
- 3. If you had a "nerve block", numbness & tingling in any part of your arm may persist for up to a day or so after surgery. This is normal. Continue to ice and work on elbow/wrist/hand range of motion.
- 4. Please call the Emergency Squad / 911 if you have sudden chest pain or severe shortness of breath.
- 5. Your post-operative office follow-up has been scheduled for you. Please call if you have any questions.
- 6. Regarding pain medicine, please call the office before 1 pm on Friday if you do not have enough pain medicines for the weekend. Most narcotic pain medicines cannot be called into your pharmacy, and the prescription must be picked up at our office.
- 7. As a group rule, we do NOT call in narcotic medicines over the weekend.

Provider Signature:	Date:
---------------------	-------