ULNAR NERVE DECOMPRESSION +/- TRANSPOSITION		CRYSTAL CLINIC ORTHOPAEDIC CENTER Post-op protocol for <i>Daniel Myer, M.D.</i>
Sling Use:		Sling for 1-3 weeks ***No PT for 1 week***
Phase I: (Passive)	Week 2-3	 Passive Range of Motion Elbow Flexion and Extension Elbow Pronation and Supination Active/Passive Hand ROM
Phase II: (Active/Assistive)	Week 4-5	 Active Assistive Range of Motion with Passive stretch to full Motion Elbow Flexion and Extension Elbow Pronation and Supination Wrist Flexion and Extension Terminal Stretching as pain allows
Phase III: (Resisted)	Week 6	 Elbow Flexion and Extension Elbow Pronation and Supination Eccentric Wrist Flexion and Extension Eccentric Wrist Pronation and Supination Standing Forward Punch Seated Rows Bicep Curls Rotator cuff strengthening PRE's for scapular strengthening
Weight Training:	Week 7	 Keep hands within eyesight, and keep elbows bent Minimize overhead activities (No Military press, pull-down behind head, or wide grip bench)
Return to activities:	Immediately	Computer, eating, holding a book, typing, writing
	2 months	Golf
	4 months	Tennis
	4 months	Contact Sports
	4 months	Throwing

